

A BIJECTION BETWEEN CERTAIN NON-CROSSING PARTITIONS AND SEQUENCES

REKHA NATARAJAN

ABSTRACT. We present a bijection between non-crossing partitions of the set $[2n+1]$ into $n+1$ blocks such that no block contains two consecutive integers, and the set of sequences $\{s_i\}_1^n$ such that $1 \leq s_i \leq i$, and if $s_i = j$, then $s_{i-r} \leq j-r$ for $1 \leq r \leq j-1$.

1. INTRODUCTION

The Catalan numbers, $C_n = \frac{1}{n+1} \binom{2n}{n}$, arise in a wide selection of combinatorial problems and applications. In his book *Enumerative Combinatorics*, Volume II, R.P. Stanley gives a list of 66 sets whose elements are counted by the Catalan numbers [5]. Later on, he has also provided an additional 26 sets in his Catalan addendum [6]. Out of this vast collection, we focus our attention on a specific type of non-crossing partitions and sequences.

Non-crossing partitions, NCP's, of the set $[n] = \{1, \dots, n\}$, where $n \in \mathbb{N}$, are partitions $\pi = \{B_1, \dots, B_k\} \in \Pi_n$, such that if $a < b < c < d$, with $a, c \in B_i$ and $b, d \in B_j$, then $i = j$ (here, Π_n denotes the set of all partitions of $[n]$.) Such partitions will always be given in lexicographic order. For instance when $n = 5$, $\pi = \{1, 5\} - \{2, 4\} - \{3\}$ is a non-crossing partition, whereas $\pi = \{1, 3\} - \{2, 4\} - \{5\}$ is a crossing partition. We choose to examine a specific class of non-crossing partitions, namely the set of non-crossing partitions of the integer $[2n+1]$ into $n+1$ blocks such that no block contains two consecutive integers, hereby referred to as 'special' partitions ([5], p. 226, Exercise 6.19(tt)). In [3], Mullin and Stanton prove that the set of special partitions of the set $[2n+1]$ is in one-to-one correspondence with the set of plane trees on $n+1$ vertices, which in turn has cardinality C_n (see [5], p. 176). For more information on these trees, see [2]. Other known bijections involving special partitions can also be found in Roselle [4].

In addition to special partitions, the Catalan numbers also count sequences $s_1, s_2, \dots, s_n$ of integers such that $1 \leq s_i \leq i$, and if $s_i = j$, then $s_{i-r} \leq j-r$ for $1 \leq r \leq j-1$ (see [5], p. 223, Exercise 6.19(z)). Stanley gives bijections between this set of sequences and the set of 312-avoiding permutations of $[n]$, as well as binary trees on n vertices ([5], p. 259, Exercise 6.19(z)).

In this paper, we present a bijection between special partitions and the sequences described above. Up until now, no bijections have been discovered between these two sets. We hope that this connection between a certain class of non-crossing partitions and sequences will shed a new light on the vast area of research encompassing permutations with some restricted patterns.

Section 2 focuses on background information relating to special partitions, while section 3 discusses the main result, which is the bijection itself.

2. DEFINITIONS AND RESULTS

We first present a few definitions and preliminary results concerning special partitions. Some of the following results may already be known, but we have not found any information regarding these in the literature. In any case, we include them here for completeness.

Definition 2.1. Let $\mathcal{P}_{[2n+1]}$ be the set of non-crossing partitions of $[2n+1]$ into $n+1$ parts such that no block contains two consecutive integers. Any partition in the set $\mathcal{P}_{[2n+1]}$ will be referred to as a ‘special’ partition.

An element $\pi \in \mathcal{P}_{[13]}$, for example, is:

$$\pi = \{1, 13\} - \{2, 4, 6, 12\} - \{3\} - \{5\} - \{7, 11\} - \{8, 10\} - \{9\}$$

We will represent π in the following manner (also known as a Puttenham diagram [1]).




FIGURE 1. $\pi = \{1, 13\} - \{2, 4, 6, 12\} - \{3\} - \{5\} - \{7, 11\} - \{8, 10\} - \{9\}$

Definition 2.2. Given a non-crossing partition $\pi = \{B_1, B_2, \dots, B_k\} \in \Pi_n$, we partition π into (disjoint) **pieces** in the following manner:

- (1) If $B_1 = 1$, then B_1 is said to be a piece of π .
- (2) If $B_1 = \{1, \dots, b_{1_w}\}$, let $B_2, \dots, B_{1+s}$ be the set of blocks in π containing the integers $2, \dots, b_{1_w} - 1$. Then the set of blocks $B_1, B_2, \dots, B_{1+s}$ is said to be a piece of π .

To find the next piece, consider $B_{1+s+1} = B_t = \{b_{1_w} + 1 = b_{t_1}, \dots, b_{t_m}\}$. Let $B_{t+1}, \dots, B_{t+i}$ be the set of blocks containing $b_{t_1} + 1, \dots, b_{t_m} - 1$. Then the set of blocks $B_t, B_{t+1}, \dots, B_{t+i}$ is also said to be a piece of π . Note that this piece could also be a singleton. Continue this process until all elements of $[n]$ have been considered.

The following examples illustrate the notion of disjoint pieces.





FIGURE 2. π made up of 3 disjoint pieces

We also need the following lemma that will be used in the proof of Theorem 2.1.

Lemma 2.1. Let $n, x_1, x_2, \dots, x_k$ be positive integers such that $n = x_1 + x_2 + \dots + x_k$. Then $\lfloor \frac{x_1}{2} \rfloor + \lfloor \frac{x_2}{2} \rfloor + \dots + \lfloor \frac{x_k}{2} \rfloor + k \geq \lfloor \frac{n}{2} \rfloor + 1$, where $\lfloor \cdot \rfloor$ denotes the floor function.

FIGURE 3. π made up of 1 piece

Proof. First note that the case $k = 1$ is trivially true. So, let $k \geq 2$ and assume that i of the x_j are even, where $i \geq 0$. Then,

$$\sum_{j=1}^k \lfloor \frac{x_j}{2} \rfloor + k = \frac{n - (k - i)}{2} + k = \frac{n}{2} + \frac{k + i}{2} \geq \lfloor \frac{n}{2} \rfloor + 1, \forall k \geq 2.$$

□

Theorem 2.1. *Let r be the number of blocks of a non-crossing partition of $[n]$ in which no block has two consecutive integers, (hereby referred to as a semi-special partition, or, SSP). Then $r \geq \lfloor \frac{n}{2} \rfloor + 1$.*

Proof. We will use induction on n .

- (1) $n = 1$. The SSP of the singleton $[1]$ contains exactly 1 block and clearly $1 \geq \lfloor \frac{1}{2} \rfloor + 1 = 1$.
- (2) $n = 2$. The SSP $\{\{1\}, \{2\}\}$ of $[2]$ contains two blocks, and again $2 \geq \lfloor \frac{2}{2} \rfloor + 1 = 2$.
- (3) Assume the claim holds for all SSP's of $k \leq n$, and let π be a SSP of $[n]$.

- (a) Let $n + 1 = 2j + 1$. By induction, π is made up of $r \geq \lfloor \frac{2j}{2} \rfloor + 1 = j + 1$ blocks. Let π' be a SSP of $[n + 1]$, with r' blocks.

If π' is obtained from π by adding $n + 1$ to an existing block, then $r' \geq j + 1 = \lfloor \frac{2j+1}{2} \rfloor + 1 = \lfloor \frac{n+1}{2} \rfloor + 1$.

Else, if $n + 1$ is added as a singleton to π , then $r' = r + 1 \geq j + 2 \geq \lfloor \frac{2j+1}{2} \rfloor + 1 = \lfloor \frac{n+1}{2} \rfloor + 1$. Thus the claim holds for $n + 1$ odd.

- (b) Let $n + 1 = 2j + 2$, and assume that π' is a SSP of $[n + 1]$ with r' blocks. We must show that $r' \geq \lfloor \frac{2j+2}{2} \rfloor + 1 = j + 2$. One sees that this inequality would fail to hold if π' is obtained by adding $n + 1$ to an existing block of π with exactly $r = \lfloor \frac{2j+1}{2} \rfloor + 1 = j + 1$ blocks. In [3], Mullin and Stanton proved that $2n + 1$ is the largest possible value of k for which there exists a SSP with $n + 1$ blocks, thus showing that $n + 1$ cannot be added to an existing block of π . However, their result follows from a theorem using face maps. Here, we give a proof relying only on elementary observations for SSP. Moreover, this proof shows that special partitions are made up of only one piece, a result which is useful in the main theorem of section 3.

We show that if π has exactly $j + 1$ blocks, then the only SSP of $n + 1$ that one can obtain from it is by adding $n + 1$ as a singleton. To prove this, we show that a SSP π of $[n] = [2j + 1]$ with exactly $j + 1$ parts has exactly one piece in the sense of Definition 2.1.

Assume that π is made up of exactly $j + 1$ blocks, which are split into 2 disjoint pieces, B and B^* (By Lemma 2.1, it suffices to consider the case where we have 2 disjoint pieces, as shown in Figure 4). Note that B and B^* are SSPs of π . Since n is odd we may assume that B




FIGURE 4. π made up of disjoint pieces

contains $2k$ integers, while B^* contains $n - 2k = 2j + 1 - 2k$ integers. By induction, B is made up of at least $k + 1$ blocks, while B^* contains at least $j - k + 1$ blocks. Together, B and B^* contain at least $j + 2$ blocks, which contradicts our assumption on π . Thus π is made up of 1 piece, and it follows that there is exactly one way to add $n + 1$ to π , namely as a singleton. Adding $n + 1$ in any other way will either violate the non-crossing condition, or will cause the two consecutive integers n and $n + 1$ to lie in the same block. Therefore, $r' = r + 1 \geq j + 2 \geq \lfloor \frac{2j+1+1}{2} \rfloor + 1 = \lfloor \frac{n+1}{2} \rfloor + 1$.

□

The following corollary is an immediate consequence of the above proof.

Corollary 2.1. *In a special partition of $[2n + 1]$, 1 and $2n + 1$ are always in the same block.*

Proof. Since a special partition consists of only 1 piece, 1 and $2n + 1$ must be in the same block. □

Corollary 2.2. *The difference between any 2 consecutive entries of any non-singleton block of a special partition is always even. That is, $b_{i_{j+1}} - b_{i_j} = 2d$.*

Proof. Consider a block $B_i \in \pi$, where π is a special partition of $[2n + 1]$. Suppose there exists a difference of $2k + 1$ between some integers b_{i_j} and $b_{i_{j+1}}$ in B_i . It follows that there are $2k$ integers trapped between b_{i_j} and $b_{i_{j+1}}$, which by Theorem 2.1 make up at least $\lfloor \frac{2k}{2} \rfloor + 1 = k + 1$ blocks. By identifying these $2k$ integers as a ‘singleton’, we can view π as a SSP of $[2n + 1 - 2k + 1]$ with at least $\lfloor \frac{2n+1-2k+1}{2} \rfloor + 1 = n - k + 2$ blocks. Thus, π is made up of at least $n - k + 2 + k + 1 - 1 = n + 2$ blocks, a contradiction. Therefore, the differences must be even. □

Definition 2.3. *Given a block B_i of size $m > 1$ in a special partition π , a subpartition is a partition formed by the set of integers between b_{i_j} and $b_{i_{j+1}}$ of B_i , $1 \leq j < m$.*

Corollary 2.3. *Subpartitions are special partitions.*

Proof. Given a special partition π of $[2n + 1]$, the subpartition on $b_{i_j} + 1, b_{i_j} + 2, \dots, b_{i_{j+1}} - 1$ is still non-crossing. Moreover, any block within the subpartition does not contain two consecutive integers. It remains to show that for a difference of $2k$ between b_{i_j} and $b_{i_{j+1}}$, the $2k - 1$ integers $b_{i_j} + 1, b_{i_j} + 2, \dots, b_{i_{j+1}} - 1$ are split into k blocks. This can be shown using the same technique as in the proof of Corollary 2.2. □

3. A BIJECTION BETWEEN SPECIAL NON-CROSSING PARTITIONS AND SEQUENCES

In this section we prove our main result, that is, we exhibit a bijection between the sets $\mathcal{P}_{[2n+1]}$ and $\mathcal{S}_n$, where $\mathcal{S}_n$ is the set of sequences $s_1, s_2, \dots, s_n$ such that the following 2 conditions hold:

- i) $1 \leq s_i \leq i$
- ii) if $s_i = j$, then $s_{i-r} \leq j - r$ for $1 \leq r \leq j - 1$

An element $s \in \mathcal{S}_6$, for instance, is:

$$s = 1 \ 2 \ 3 \ 1 \ 1 \ 6$$

The following observations about sequences in $\mathcal{S}_n$ will be useful: The only integer s_i such that $s_i = s_{i+1}$ is $s_i = 1$.

Lemma 3.1. *Any sequence $s \in \mathcal{S}_n$ can be obtained from a sequence of n 1s using the following algorithm.*

- (1) Start with the two sequences T and g , where $T = 1 \ 1 \ \dots \ 1$ (a sequence of n 1's), and $g = 1 \ 2 \ \dots \ n$.
- (2) Build successive pairs of sequences $\{T^1, g^1\}, \{T^2, g^2\}, \dots, \{T^n, g^n\}$ in the following manner:
 - (a) Pick an integer $m_1 \leq n = g_n$. Let $T^1 = T_1^1 \ T_2^1 \ \dots \ T_n^1$, where $T_i^1 = T_i$ for all i such that $i < n$, and $T_n^1 = m_1$.
Let $g^1 = g_1^1 \ g_2^1 \ \dots \ g_n^1$, where $g_i^1 = g_i$ for all i such that $i \leq n - m_1$, and $g_{n-m_1+j}^1 = j$ for all j such that $1 \leq j \leq m_1$. Note that choosing $m_1 = 1$ is also a possibility, in which case, $g^1 = 1 \ 2 \ \dots \ (n-1) \ 1$, and $T^1 = T$.
 - (b) Build T^2 from T^1 by replacing the $(n-1)^{th}$ term of T^1 , T_{n-1}^1 , with any integer $m_2 \leq g_{n-1}^1$.
Let $g^2 = 1 \ 2 \ \dots \ n - m_1 \ 1 \ 2 \ \dots \ (m_1 - m_2 - 1) \ 1 \ 2 \ \dots \ m_2 \ m_1$. Note that if $m_2 < m_1$, then $m_1 - m_2 - 1 \geq 0$. In the case $m_2 = m_1 + 1$, we simply have that $g^2 = g^1$.
 - (c) Continue building such pairs of sequences $\{T^i, g^i\}$ where for all i , T^i is obtained from T^{i-1} by replacing the one in position $(n-i+1)$ with any integer $m_i \leq g_{n-i+1}^{i-1}$. The sequence g^i is built from the sequence g^{i-1} in the manner described earlier. Notice that for all i , $T^i, g^i \in \mathcal{S}_n$.
One clearly sees that any sequence $s \in \mathcal{S}_n$ can be obtained in the manner described above, and that indeed the only sequences obtained from this algorithm belong to $\mathcal{S}_n$.

The following example illustrates this lemma.

Example. Pairs of sequences obtained from the algorithm above, where $n = 8$.

$$\begin{array}{ll}
T : 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 & g : 1 \ 2 \ 3 \ 4 \ 5 \ 6 \ 7 \ 8 \\
T^1 : 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 4 & g^1 : 1 \ 2 \ 3 \ 4 \ 1 \ 2 \ 3 \ 4 \\
T^2 : 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 4 & g^2 : 1 \ 2 \ 3 \ 4 \ 1 \ 2 \ 1 \ 4 \\
T^3 : 1 \ 1 \ 1 \ 1 \ 1 \ 2 \ 1 \ 4 & g^3 : 1 \ 2 \ 3 \ 4 \ 1 \ 2 \ 1 \ 4 \\
T^4 : 1 \ 1 \ 1 \ 1 \ 1 \ 2 \ 1 \ 4 & g^4 : 1 \ 2 \ 3 \ 4 \ 1 \ 2 \ 1 \ 4 \\
T^5 : 1 \ 1 \ 1 \ 4 \ 1 \ 2 \ 1 \ 4 & g^5 : 1 \ 2 \ 3 \ 4 \ 1 \ 2 \ 1 \ 4
\end{array}$$

Given $\pi \in \mathcal{P}_{[2n+1]}$, where $B_i = \{b_{i_1}, b_{i_2}, \dots, b_{i_m}\}$, we construct a sequence $d_1, d_2, \dots, d_{2n+1}$ of differences in the following manner:

- (1) Let $i = b_{j_k} \in B_j$. Then let
 - a) $d_i = b_{j_{k+1}} - i$ if $|B_j| > k$.
 - b) $d_i = 0$ if $|B_j| = k$.

Since a special partition $\pi \in \mathcal{P}_{[2n+1]}$ is made up of $n+1$ blocks, there are exactly $n+1$ differences that are zero. Thus, there are $2n+1 - (n+1) = n$ non-zero differences.

- (2) Let $\{d_{i_k}\}_1^n$ be the subsequence of non-zero integers, and form a new sequence $\{a_j\}_1^n$ such that $a_j = (d_{i_{n-j+1}})/2$.

Define a map $f : \mathcal{P}_{[2n+1]} \mapsto [n-1]^n$ by $f(\pi) = (a_1, a_2, \dots, a_n)$, where the sequence $(a_1, a_2, \dots, a_n)$ is obtained using steps 1 and 2 as described above.

Example. $\pi = \{1, 13\} - \{2, 4, 6, 12\} - \{3\} - \{5\} - \{7, 11\} - \{8, 10\} - \{9\}$

The sequence of differences as obtained from step 1 is:

$$d_1 = 12, d_2 = 2, d_3 = 0, d_4 = 2, d_5 = 0, d_6 = 6, \\ d_7 = 4, d_8 = 2, d_9 = 0, d_{10} = 0, d_{11} = 0, d_{12} = 0, d_{13} = 0$$

The subsequence of non-zero integers is:

$$\{d_{i_k}\}_1^6 = : \{12 \quad 2 \quad 2 \quad 6 \quad 4 \quad 2\}$$

The sequence after reversing and dividing by two is:

$$\{a_j\}_1^6 = : \{1 \quad 2 \quad 3 \quad 1 \quad 1 \quad 6\}$$

It is easy to check that this sequence is an element of $\mathcal{S}_6$. In Corollary 2.2, we showed that all differences d_i are even. Thus, the sequence $\{a_j\}$ obtained from f is indeed a sequence of integers.

Theorem 3.1. The mapping f is a bijection between $\mathcal{P}_{[2n+1]}$ and $\mathcal{S}_n$.

Proof. First, we show that f is well-defined. The maximum value of d_1 is $2n+1-1 = 2n$. But $(d_1)/2$ corresponds to a_n , so $a_n \leq n$. Similarly, the maximum value of d_{i_k} is $2n - 2k + 2$. Thus,

$$a_j = d_{i_{n-j+1}}/2 \leq (2n - 2(n-j+1) + 2)/2 = 2j/2 = j,$$

showing that the first condition for being a sequence in the set $\mathcal{S}_n$ holds. By Corollary 2.3, any difference of $2j$ between two integers b_{i_k} and $b_{i_{k+1}}$ of a block gives rise to a subpartition with $j-1$ differences. The maximum values of these differences are $2j-2, 2j-4, \dots, 2j-2(j-1)$. Therefore, $a_{j-r} \leq j-r$, $1 \leq r \leq j-1$, and condition 2 holds. Thus, f is well-defined.

To show that f is onto, given a sequence $s = s_1 s_2 \dots s_n \in \mathcal{S}_n$, we build its corresponding special partition in the following manner:

Associate the sequence of n 1s with the Puttenham diagram $\mathcal{D}_1$ below:




FIGURE 5. $\mathcal{D}_1$: Puttenham diagram for a sequence with n 1's, $n = 6$

Note that $\mathcal{D}_1$ contains $n + 1$ blocks and is clearly special. We will construct a sequence of special partitions from $\mathcal{D}_1$ as follows. First, in order to obtain the sequence s from the algorithm described in Lemma 3.1, observe that $T_n^1 = s_n$. Now, create a corresponding Puttenham diagram $\mathcal{D}_2$ by replacing the 1st arc of $\mathcal{D}_1$, $(1, 3)$, with the arc stretching from 1 to $1 + 2s_n$, and by shifting $s_n - 1$ arcs of length two underneath it. In other words, the arcs $(3, 5), (5, 7), (7, 9), \dots$, are replaced by $(2, 4), (4, 6), (6, 8)$ and so on. Note that if $s_n = 1$, then $\mathcal{D}_2 = \mathcal{D}_1$. Observe that $\mathcal{D}_2$ contains $n + 1$ blocks, and is clearly special. Moreover, $T^1 = 1 \ 1 \ \dots \ s_n$ and $g^1 = 1 \ 1 \ \dots \ (n - s_n) \ 1 \ 2 \ \dots \ n$, which means that for the next $s_n - 1$ steps, we will be transforming $\mathcal{D}_2$ by working ‘underneath’ the arc $(1, 1 + 2s_n)$.

Next, $T^2 = 1 \ 1 \ \dots \ s_{n-1} \ s_n$, and we obtain $\mathcal{D}_3$ from $\mathcal{D}_2$ by replacing the second arc of $\mathcal{D}_2$, $(2, 4)$, with the arc $(2, 2 + 2s_{n-1})$ and once again shifting arcs of length two underneath as before. Again, if $s_{n-1} = 1$, then $\mathcal{D}_3 = \mathcal{D}_2$.

Continue creating each $\mathcal{D}_{i+1}$ in the following manner: Stretch the i th arc of $\mathcal{D}_i$ by $2(T_{n-i+1}^i) = 2s_{n-i+1}$, and shift $T_{n-i+1}^i - 1 = s_{n-i+1} - 1$ arcs ‘underneath’ it. Notice that after $s_n - 1$ steps, we repeat the process of stretching and shifting arcs at the integer $1 + 2s_n$, and so on.

The reason that we are able to shift arcs in the manner described above is because each T^j has a corresponding governing sequence g^j whose $(n - j + 1)^{th}$ entry helps us identify the exact number of arcs of length two that will fit “underneath” the stretched arc in $\mathcal{D}_{j+1}$. Also, notice that at each step, $\mathcal{D}_{j+1}$ is made up of exactly one piece.

Clearly, each time arcs are stretched and shifted, the number of blocks in the diagram $\mathcal{D}_{j+1}$ is preserved, as are the conditions of being a special partition. Note that subpartitions of each $\mathcal{D}_{j+1}$ are special partitions as well. After n steps (or less), this method of construction gives us an element in $\mathcal{P}_{[2n+1]}$ that corresponds to s and we are done. The figure below illustrates how to construct the special partition corresponding to $s = 1 \ 1 \ 1 \ 4 \ 1 \ 2 \ 1 \ 4$, where $n = 8$. Notice here that $T^5 = T^6 = T^7 = T^8$, and therefore $\mathcal{D}_6 = \mathcal{D}_7 = \mathcal{D}_8 = \mathcal{D}_9$.




FIGURE 6. Constructing the special partition corresponding to $s = 1\ 1\ 1\ 4\ 1\ 2\ 1\ 4$.

□

4. ACKNOWLEDGMENTS

Sincere thanks are due to Hélène Barcelo for helpful discussions and suggestions.

REFERENCES

- [1] H.W. Becker, *Planar Rhyme Schemes*, Bulletin of the American Mathematical Society. 1952.
- [2] D.A. Klarner, *Correspondences between Plane Trees and Binary Sequences*, Journal of Combinatorial Theory, vol. 9, 1970, pg. 401-411.
- [3] R.C. Mullin and R.G. Stanton, *A Map-Theoretic Approach to Davenport-Schinzel Sequences*, Pacific Journal of Mathematics, vol. 40, No. 1, 1972.
- [4] D.P. Roselle, *An Algorithmic Approach to Davenport-Schinzel Sequences*, Utilitas Math, vol. 6, 1974, 91-93.
- [5] Richard P. Stanley, *Enumerative Combinatorics, vol. II*, Cambridge University Press, 1999.
- [6] Richard P. Stanley, *Catalan Addendum* August 19, 2002. www-math.mit.edu/~rstan/ec.